

Table I. *Relative level of KGFR expression in human carcinomas.*

Cancer type	Breast	Ovary	Uterus	Cervix	Vulva	Prostrate	Testes	Thyroid	Skin
KGFR ⁺	(6/10)	(10/10)	(7/10)	(3/10)	(1/5)	(3/4)	(8/10)	(1/10)	(0/10)
KGFR Ratio*	4.56	1.07	2.99	0.66	0.33	1.18	1.51	0.31	0.63
	5.79	2.94	2.66	0.85	0.37	0.90	1.76	0.32	0.27
	0.57	4.10	1.21	0.63	0.53	2.31	1.47	0.50	0.40
	1.65	1.70	0.79	0.72	0.24	1.74	0.35	0.95	0.74
	0.63	4.94	1.07	1.85	1.65		1.47	0.32	0.21
	0.59	1.92	0.92	1.61			3.48	1.15	0.50
	1.06	3.49	2.55	0.26			0.68	1.00	0.39
	0.49	1.30	0.61	0.61			2.76	0.28	0.46
	1.32	2.28	4.54	0.55			2.34	0.47	0.34
1.29	1.35	3.69	2.37			2.75	0.86	0.59	
Cancer type	Stomach	Small Intestine	Colon	Rectum	Pancreas	Liver	Lung	Kidney	Bladder
KGFR ⁺	(6/10)	(5/7)	(1/10)	(4/10)	(2/7)	(0/3)	(6/10)	(1/10)	(2/5)
KGFR Ratio*	1.46	0.96	0.48	0.84	0.72	0.25	1.34	0.34	0.88
	0.88	1.27	0.80	1.11	0.54	0.58	0.77	0.71	0.58
	34.91	0.67	1.24	3.58	1.30	0.23	0.75	0.47	1.17
	0.86	1.40	0.77	0.90	0.45		3.08	0.30	0.89
	0.74	1.32	0.84	0.42	1.05		0.64	0.17	2.59
	1.31	2.33	0.90	1.98	0.25	Trachea (2/3)	2.23	0.19	
	4.35	1.24	0.94	0.78	0.45		1.04	0.58	
	41.17		0.40	0.91		1.76	4.65	0.23	
	1.42		0.87	1.00		0.87	0.38	0.94	
0.82		0.55	1.07		1.36	2.65	1.01		

*Fraction of tumors with KGFR over-expressed relative to the patient-paired normal tissue within the cancer type subset. *Ratio of KGFR hybridization in tumor samples relative to the patient-paired normal tissue.