

Table II. Comparative CPs of gene expression in isolated colonocytes and stool samples from normal individuals and cancer patients.

Group	Isolated colonocytes									Stool samples								
	HPRT	IGF2	FLNA	TGFβ _{high}	CKS2	CSE1L	DPEP1	KLKI0	CXCL3	HPRT	IGF2	FLNA	TGFβ _{high}	CKS2	CSE1L	DPEP1	KLKI0	CXCL3
NegCont ^a	37.36	36.48	37.79	37.83	37.55	36.49	37.57	36.49	36.33	37.32	37.07	37.55	36.69	36.88	37.22	36.98	37.36	37.11
NorC/S1	36.66	37.46	36.31	36.09	37.14	37.81	36.79	37.22	37.43	37.22	36.58	36.94	37.35	37.07	36.97	37.58	36.52	37.08
NorC/S2	37.48	37.29	37.03	37.42	37.77	36.92	37.25	37.43	36.87	36.99	37.22	37.11	36.79	37.14	37.61	36.83	37.18	37.36
C/S0-13	37.29	23.55	25.01	27.56	29.89	31.68	32.67	33.82	35.17	37.16	23.21	24.78	27.22	30.01	31.15	32.22	33.69	34.99
C/S0-14	36.79	24.01	25.11	27.87	30.03	31.58	32.91	33.79	35.01	37.52	23.69	25.09	27.59	29.37	30.99	33.02	33.83	35.15
C2/S25	37.22	22.84	23.59	25.17	26.55	27.99	29.15	31.02	31.68	37.11	22.92	23.66	25.07	26.14	28.02	29.07	30.89	32.33
C2/S26	36.79	22.41	23.36	25.48	26.89	27.69	29.09	30.87	31.99	37.07	23.04	23.57	25.61	26.55	27.89	29.18	30.71	31.87
C2/S27	37.37	22.15	23.75	25.08	26.33	27.45	29.57	30.51	31.44	36.78	22.45	23.49	25.45	26.11	27.61	29.26	30.43	32.04
C3/S38	37.26	19.45	21.36	24.68	25.78	26.89	28.58	29.41	30.99	37.11	19.23	21.42	24.52	25.61	26.71	28.42	29.13	30.84
C3/S39	36.15	19.51	21.11	23.98	25.43	26.76	28.43	29.32	31.03	37.82	18.98	21.33	24.09	25.33	26.19	28.21	29.45	30.88
C3/S310	37.09	18.98	21.79	23.35	25.13	26.35	28.27	29.67	31.27	37.31	18.75	21.44	23.67	25.63	26.69	29.01	29.59	31.46

^aNo DNA added to reaction (negative control). This table was obtained from 10 individuals: 2 non-cancerous controls and 8 patients with colon cancer. NorC/S1 and NorC/S2=Isolated colonocytes/Stool from normal individuals 1 and 2; C₀₋₁/S₀₋₁ 4 and C₀₋₁/S₀₋₁ 5=Isolated colonocytes/Stool from patients 3 and 4 with adenomatous colon polyp ≥1 cm (stage 0-1); C₂/S₂ 5 to S/C₂7=Isolated colonocytes/Stool from colon carcinoma patients 5 to 7 (stage 2); C₃/S₃ 8 to 10=Isolated colonocytes/Stool from colon carcinoma patients 8 to 10 (stage 3). All samples were preserved overnight at 4 °C in S.T.A.R[®] preservative before RNA extraction.